

FIFRA Section 24(c) Special Local Need



DOC ID# 592637

DINOTEFURAN	GROUP	4A	INSECTICIDE
-------------	-------	-----------	-------------

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW YORK

ACCEPTED
FOR REGISTRATION

Mar 14, 2024

SAFARI® 20 SG INSECTICIDE

**Basal Trunk Spray for Control of Hemlock Woolly Adelgid
and Elongate Hemlock Scale on Hemlocks Only**

Classified for
“RESTRICTED USE”
in New York State
under 6NYCRR Part 326

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

EPA Reg. No. 86203-11-59639
EPA SLN No. NY-120009

This label for Safari® 20 SG Insecticide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2028.

Active Ingredient:	By Wt
Dinotefuran, [N-methyl-N'-nitro-N''-(tetrahydro-3-furanyl)methyl]guanidine]	20%
Other Ingredients	80%
Total	100%

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	PESTS	APPLICATION RATE FOR BASAL TRUNK SPRAY
Eastern Hemlock (<i>Tsuga canadensis</i>)	Hemlock Woolly Adelgid (<i>Adelges tsugae</i>)	9.0 oz <i>Safari</i> 20 SG per gallon of water (=1.8 oz ai per gallon of water)
Carolina Hemlock (<i>Tsuga caroliniana</i>)	Elongate Hemlock Scale (<i>Fionnia externa</i>)	When applied just to point of runoff, one gallon of spray solution should treat approximately 65-85 inches of cumulative trunk diameter (=1.5-2.0 fl oz spray solution per inch of trunk diameter).

The active ingredient in *Safari* 20 SG is a highly systemic product that will move through the bark into the xylem and then be transported upward to where hemlock woolly adelgid and elongate hemlock scale feed. In actively growing hemlocks, control may be evident within 2 weeks after application.

APPLICATION INSTRUCTIONS:

- Apply when hemlocks are actively growing and transporting water in xylem (March-November).
- Apply with a low-pressure sprayer operated at maximum of 20 PSI to avoid damage to the trees, splash-back and drift.
- Spray solution on bark over entire circumference of tree trunk from 4” – 54” above the soil surface. Wet bark with spray solution just to the point of saturation and run-off. During application, do not allow spray solution to run off trunk on to soil surface.
- Do not apply directly to soil.
- Do not directly apply to any plant species other than Eastern or Carolina hemlocks.
- Do not apply to rain saturated bark, during rainfall, or if significant rainfall is expected within 24 hours after application.
- Do not make more than 2 applications per hemlock tree per year
- Do not apply more than 2.7 lb of *Safari* 20 SG per acre per year (=0.54 lb ai per acre per year).

GENERAL INFORMATION

Safari 20 SG may be applied as a low-pressure basal trunk spray to hemlock trees growing in outdoor sites including apartment complexes, athletic fields, cemeteries, golf courses, lawns and ornamental landscapes around residences and businesses, parks, military bases and other government institutions, office complexes, playgrounds, and in private, city, state and federal parks, recreation areas and forests

APPLICATION EQUIPMENT

Safari 20 SG Insecticide can be applied using many different types of low-pressure sprayers. Before applying *Safari* 20 SG Insecticide, start with a clean, well maintained sprayer. Use a spray nozzle that produces medium-sized droplets.

MIXING INSTRUCTIONS

1. Fill clean spray tank to 2/3 of desired level with clean water.
2. Add *Safari* 20 SG Insecticide, agitate until product has dissolved and then fill spray tank to desired level with clean water.
3. Continue agitation until all spray solution has been applied.

Follow all applicable directions, restrictions and precautions on the EPA-registered label. This labeling must be in the possession of the user at the time of pesticide application.

FIFRA § 24(c) Registrant: Valent U.S.A. LLC
P.O. Box 5075
San Ramon, CA 94583

Safari is a registered trademark of Valent U.S.A. LLC.